

Rigidity Sensing at the Leading Edge through $\alpha_v\beta_3$ Integrins and RPTP α

Guoying Jiang, Angela H. Huang, Yunfei Cai, Monica Tanase, and Michael P. Sheetz

Department of Biological Sciences, Columbia University, New York, New York 11027

ABSTRACT Cells require optimal substrate stiffness for normal function and differentiation. The mechanisms for sensing matrix rigidity and durotaxis, however, are not clear. Here we showed that control, Shp2^{-/-}, integrin β_1 ^{-/-}, and talin1^{-/-} cell lines all spread to a threefold greater area on fibronectin (FN)-coated rigid polyacrylamide surfaces than soft. In contrast, RPTP α ^{-/-} cells spread to the same area irrespective of rigidity on FN surfaces but spread 3 × greater on rigid collagen IV-coated surfaces than soft. RPTP α and $\alpha_v\beta_3$ integrins were shown previously to be colocalized at leading edges and antibodies to $\alpha_v\beta_3$ blocked FN rigidity sensing. When FN beads were held with a rigid laser trap at the leading edge, stronger bonds to the cytoskeleton formed than when held with a soft trap; whereas back from the leading edge and in RPTP α ^{-/-} cells, weaker bonds were formed with both rigid and soft laser traps. From the rigidity of the trap, we calculate that a force of 10 pN generated in 1 s is sufficient to activate the rigidity response. We suggest that RPTP α and $\alpha_v\beta_3$ at the leading edge are critical elements for sensing FN matrix rigidity possibly through SFK activation at the edge and downstream signaling.

INTRODUCTION

The stiffness of extracellular matrix (ECM) dramatically affects many cellular processes, such as motility (1,2), phagocytosis (3), and cell differentiation (4). However, different cell types prefer different matrix rigidities (5). Normal fibroblasts do not spread fully, form extensive focal contacts, or grow on soft substrates (2,6). Endothelial cells from human umbilical vein are more spread and have larger lumens and less branching on stiffer collagen gels (7). Myocytes only formed myotubes with striations on gels of intermediate stiffness, and changes in myoblast behavior were related to differences in stiffness of healthy and diseased muscle tissue (5,8). On the other hand, hepatocytes maintain a differentiated phenotype only on soft materials, and changes in mechanical properties during liver disease may be partially responsible for the deterioration of hepatocyte networks (5,9). Oncogenically transformed cells can grow on gelatin-coated materials softer than 100 Pa, whereas nontransformed cells cannot survive (10,11). Primary neuron cultures from mouse spinal cords branched more frequently on soft substrates (12). Thus, the rigidity response can contribute to differentiation of cells within a particular tissue (5) and is a major factor in cancers and other disease states as well (13,14).

In fibroblast cells, matrix-integrin interactions on the active lamellipodia cause indirect attachment of integrins to actin filaments (15). As the filaments are moved rearward by myosin motors, they generate force on the matrix when it resists movement. Force increases rapidly with small movements when the matrix is rigid or more slowly with larger movements when the matrix is soft (16). Fibroblasts sense substrate rigidity and move toward rigid areas both in three

dimensions (4,17) and in two dimensions, a phenomenon defined as durotaxis (1), by an unknown process(es). Generation of periodic contractions in extending lamellipodia appears to be linked to the mechanical probing of the ECM rigidity by the cell (18). However, to date, no specific molecule has been shown to be the sensor of the fibronectin (FN) matrix rigidity although periodic rows of β_3 integrin clusters were observed in spreading cells (18).

$\alpha_v\beta_3$ integrin has important roles in the migration and invasion of melanoma cells (19,20), vascular endothelial cells (21), and primary tumor growth and metastasis in vivo (20,21). It forms a complex with RPTP α , a receptor-like protein tyrosine phosphatase, at the leading edge early in spreading (22). RPTP α was identified previously as a component involved in the cellular response to force (22). Gene inactivation of RPTP α delays spreading on FN, impairs activation of Src family kinases (SFK) (23,24), and compromises correct positioning of pyramidal neurons during development of mouse hippocampus (25). Those studies identified RPTP α as a key component for proper radial neuronal migration (25). In this study, we investigated the function of $\alpha_v\beta_3$ integrin and RPTP α in FN matrix rigidity sensing by measuring the cell spread area on different stiffness polyacrylamide gel surfaces and the strength of cytoskeleton bonding to FN-coated beads with laser traps of different stiffness. Our data indicate that rigidity sensing involves position-dependent changes in force on the RPTP α / $\alpha_v\beta_3$ complex at the leading edge.

METHODS AND MATERIALS

Cell culture and materials

Mouse fibroblast cells (FAK^{+/+}, FAK^{-/-}, RPTP α ^{+/+}, RPTP α ^{-/-}, Shp2^{+/+}, Shp2^{-/-}, β_1 ^{-/-}, Talin1^{-/-}) were maintained at 37°C and 5% CO₂ in Dulbecco's Modified Eagle's medium (Invitrogen, Carlsbad, CA) supplemented with 10% fetal bovine serum (Invitrogen), 2 mM L-glutamine, 50 μ g/ml streptomycin, and 50 units/ml penicillin. Anti- $\alpha_v\beta_3$ (clone LM609,

Submitted August 13, 2005, and accepted for publication October 26, 2005.

Address reprint requests to Michael P. Sheetz, Dept. of Biological Sciences, Columbia University, 1212 Amsterdam Ave., New York, NY 11027. Tel.: 212-854-4857; Fax: 212-854-6399; E-mail: ms2001@columbia.edu.

© 2006 by the Biophysical Society

0006-3495/06/03/1804/06 \$2.00

doi: 10.1529/biophysj.105.072462

Chemicon, Temecula, CA) antibody was included at optimal concentration in the integrin binding inhibition experiments.

Cell spreading on polyacrylamide substrates and microscopy

The polyacrylamide substrates were prepared and coated with FN and collagen IV as described previously (2). The flexibility of the substrate was manipulated by maintaining the total acrylamide concentration at 8% while varying the bis-acrylamide components between 0.4% (rigid surface) and 0.03% (soft surface). The Young's modulus of the polyacrylamide substrates was measured and calculated as described by Pelham et al. (2). The uniformity of FN coating on the substrate surface was examined by coating the gels with Alexa 568 labeled FN and observed by immunofluorescence microscopy. Experiments were performed 15 h after the cells were plated on the polyacrylamide gel at a low density. Phase contrast images were recorded with a cooled charge-coupled device camera attached to an Olympus IX81 equipped with a 10 \times objective. The spread area of individual cells was quantified with Image J software. At least 50 cells were counted for each cell line under each condition.

Breaking events assay

Cells were plated on acid-washed, silane-treated coverslips coated with laminin (40 $\mu\text{g}/\text{ml}$). With an optical-gradient laser trap as described previously (26), FN-coated beads were held at the leading edge or 2 μm away from the leading edge of the lamellipodia for 3 s, then released by turning off the laser power. If the bead bound to the cell membrane, the laser was turned on again. The movement of the bead restrained by the laser trap was recorded until the bead was finally pulled out of the trap. Breaking events were defined by the rapid movement of beads to the center of the laser trap after the cell started to move the bead rearward.

RESULTS AND DISCUSSION

Rigidity-dependent spreading requires RPTP α

To determine which protein(s) may be involved in the rigidity-sensing process, we screened a number of knockout cell lines for their ability to spread on rigid versus soft polyacrylamide substrates coated with FN (see Fig. 1 A). As shown in Fig. 1 B, the spread areas of wild-type cells (RPTP α ^{+/+}, FAK^{+/+}, Shp2^{+/+}) were two- to threefold greater on rigid than on soft surfaces. Similarly, Shp2^{-/-}, talin1^{-/-}, and integrin β 1^{-/-} cells spread more on rigid than on soft surfaces.

In contrast, FAK^{-/-} cells had only a very small contact area on rigid surfaces but spread to the same area as controls on soft polyacrylamide surfaces. Since myosin or Rho-kinase (ROCK) inhibition caused FAK^{-/-} cell contact areas to increase to control cell areas on glass (27), we believed that hypercontraction produced the small contact area. In our assay of FAK^{-/-} cell spreading on rigid FN polyacrylamide gels, ROCK inhibition caused an increase in the contact area (data not shown). Consequently, we believed that rigidity was sensed in the FAK^{-/-} cells and the normal increase in contraction on rigid surfaces was accentuated in the absence of FAK.

We found one mutant cell line, RPTP α ^{-/-} cells, that spread to the same area on soft and rigid surfaces ($p < 0.21$, t -test)




FIGURE 1 Screening of FN rigidity sensing response of different cell lines. (A) Control cells were spread on rigid versus soft FN-coated polyacrylamide substrates. The spread area of individual cells was measured with Image J. Scale bar, 40 μm . (B) Spread area of different cell lines on rigid and soft FN substrates. Results shown are the mean \pm SD of at least 50 cells. Asterisks indicate significant statistical difference between the rigid and soft surface, $p < 0.01$.

(Figs. 1 B and 3 B). When cells were cultured on FN-coated surfaces with varying stiffness (Fig. 2 B) by changing the concentration of bis-acrylamide (0.03%, 0.08%, 0.2%, and 0.4%), RPTP α ^{+/+} cells spread to larger areas as the matrix rigidity increased, whereas RPTP α ^{-/-} cells did not show the stiffness-dependent spreading behavior (Fig. 2 C). When the level of RPTP α in wild-type cells was reduced by siRNA (Fig. 2 A), cells spread to an even smaller area than RPTP α ^{-/-} cells and were unable to sense the matrix rigidity changes (Fig. 2 C). Thus, we suggest that RPTP α is critical for the matrix rigidity sensing process.

$\alpha_v\beta_3$ integrin involvement in the rigidity sensing process

Because RPTP α formed a complex with $\alpha_v\beta_3$ at the leading edge (22), we incubated control fibroblasts with anti- $\alpha_v\beta_3$ monoclonal antibody LM609 at 10 $\mu\text{g}/\text{ml}$ (28,29) during cell spreading. As shown in Fig. 3 A, RPTP α ^{+/+} cells treated with LM609 showed no difference in spread area on rigid versus soft FN polyacrylamide surfaces ($p < 0.1$, t -test), which indicated that the cells lost their ability to sense FN rigidity when the binding of $\alpha_v\beta_3$ to FN was blocked. Treatment of RPTP α ^{-/-} cells with LM609 had no effects on the spread area ($p < 0.45$, t -test). Thus, $\alpha_v\beta_3$ integrin was an important component for FN rigidity sensing at the leading




FIGURE 2 Rigidity sensing of FN matrix is RPTP α dependent. (A) Western blot showing the expression of RPTP α in control (CT) and knockdown (A1 and A2) cells. (B) Mechanical properties of polyacrylamide substrate. The Young's modulus (N/m²) of polyacrylamide gels with a range of bis-acrylamide (0.03%, 0.08%, 0.2%, and 0.4%) to acrylamide (8%) was measured as described by Pelham et al. (2). (C) The spread areas of RPTP $\alpha^{+/+}$, RPTP $\alpha^{-/-}$, and RPTP α knockdown (A2) cells on polyacrylamide substrates of different rigidities. Results shown are the mean \pm SD of at least 50 cells.

edge during early spreading, possibly through an interaction with RPTP α .

To determine if RPTP $\alpha^{-/-}$ cells could sense matrix rigidity through other integrins that did not bind to RPTP α (22), we plated cells on rigid versus soft polyacrylamide substrate coated with collagen IV, which binds to the $\alpha_1\beta_1$ integrin (30). Both control and RPTP $\alpha^{-/-}$ cells spread to threefold greater area on the rigid collagen IV surfaces (Fig. 3 C). Thus, it was clear that $\alpha_1\beta_1$ integrins can participate in rigidity sensing through a mechanism that does not involve RPTP α .

Matrix rigidity sensing at the leading edge

Because the two membrane proteins ($\alpha_v\beta_3$ and RPTP α) implicated in rigidity sensing were at the leading edge (22), we compared rigidity sensing at the edge and two microns back from the edge using the laser trap microscope (Fig. 4 A). The




FIGURE 3 Rigidity sensing of FN matrix is dependent on $\alpha_v\beta_3$ integrins and RPTP α , whereas rigidity sensing on collagen IV matrix is RPTP α independent. (A) Spread area of RPTP $\alpha^{+/+}$ and RPTP $\alpha^{-/-}$ cells plated on FN-coated polyacrylamide substrate with or without 10 $\mu\text{g}/\text{ml}$ LM609. (B) The morphologies of RPTP $\alpha^{-/-}$ cells on FN and Collagen IV (Col)-coated substrates. (C) Spread area of RPTP $\alpha^{+/+}$ and RPTP $\alpha^{-/-}$ cells plated on collagen IV-coated polyacrylamide substrates. Results shown are the mean \pm SD of at least 50 cells. Asterisks indicate a significant statistical difference between the rigid and soft surface, $p < 0.01$. Scale bar, 40 μm .

rigidity of beads held in the laser trap was directly proportional to the laser power. There were two cellular responses to rigid surfaces: 1), activation of leading edge extension (18), and 2), increased strengthening of bonds between




FIGURE 4 Rigidity sensing is related to the strength of FN bead linkages to the cytoskeleton and is RPTP α dependent. (A) Schematic graph showing the position of rigid or soft trap on the lamellipodia of a spreading cell. (B) Breaking percentage of FN beads placed on the lamellipodia of RPTP $\alpha^{+/+}$ cells. (C) Breaking percentage of FN beads placed on the lamellipodia of RPTP $\alpha^{-/-}$ cells. Results shown are the mean \pm SD of three independent experiments. Asterisks indicate significant statistical difference between the two groups, $p < 0.01$.

integrins and the cytoskeleton with increased recruitment of focal complex proteins (22). Increased extension was manifested as an increase in spread area (as above); however, increased strengthening of bonds could be measured by increased resistance of beads to movement by the laser trap. To quantify bond strength, we measured the frequency of breaking of bonds between FN beads and the actin cytoskeleton as the cell moved the beads toward the nucleus at the rate of ~ 60 nm/s. With soft (0.02 pN/nm) laser trap, there was nearly twice the number of breaking events than with

rigid (0.18 pN/nm) trap (Fig. 4 B), which indicated that the rigid trap caused stronger bonds to the cytoskeleton to form at the leading edge. When beads were placed 2 microns back from the edge, there was the same number of breaking events for both rigid and soft laser trap, matching the weaker bonding with the soft trap at the edge. Thus, the rigid trap was only sensed at the leading edge, where it increased the strengthening of FN-cytoskeleton linkages.

An alternative explanation for the increased bonding to the cytoskeleton with the increased laser trap rigidity was that the increased light intensity caused photo-induced cross-linking. To control for possible photodamage, we made the soft trap appear rigid by rapidly shifting the stage 500 nm to the point of highest force in the trap, causing a rapid increase in force. When the 0.02 pN/nm laser trap was moved 500 nm from the cell center to generate higher force instantaneously, we found fewer bond-breaking events (data not shown). This indicated that the soft trap could mimic rigid trap by a rapid increase in the force on the bead, which caused stronger bonds to form between the beads and the cytoskeleton without laser photodamage.

If RPTP α was involved in the bead response to laser trap rigidity, then RPTP $\alpha^{-/-}$ cells should not show rigidity dependence in the frequency of breaking events even at the edge. Indeed, the frequency of breaking events was the same for soft and rigid laser traps at the edge (Fig. 4 C), and the high fraction of breaking events was similar to the frequency when beads were placed 2 microns back from the edge. Thus, we found that the edge response to laser traps rigidity was dependent upon RPTP α .

Rigidity sensing downstream of $\alpha_v\beta_3$ and RPTP α

In this study we demonstrate that the membrane proteins RPTP α and $\alpha_v\beta_3$ are involved in the sensing of FN matrix rigidity at the leading edge during early spreading. Further steps in the rigidity-sensing pathway are in question. Previous experiments have demonstrated that $\alpha_v\beta_3$ integrins and RPTP α are involved in the activation of SFKs during early spreading (22). Either the absence of RPTP α or inhibition of FN binding to $\alpha_v\beta_3$ (with an antibody or an inhibitory peptide, Gpen) significantly reduces SFK activation (22), which parallels the effects on rigidity sensing. In contrast, talin1 $^{-/-}$ cells, which are defective in force-dependent reinforcement of FN bead-cytoskeleton bonds but have normal SFK activation (31), display normal rigidity sensing behavior (Fig. 1 B). Thus, we suggest that SFK activation is critical in the FN rigidity sensing process.

Previous studies indicated that the sensing of rigid substrates involves either the stimulation of a tyrosine kinase or inhibition of a tyrosine phosphatase (2). With more rigid substrates, there is greater tyrosine phosphorylation in NIH3T3 fibroblasts; and cells on soft substrates spread to the area of cells on rigid substrates after treatment with PAO, a phosphatase inhibitor (2). Our data suggest more specifically that the




FIGURE 5 Model of FN matrix rigidity sensing. On a rigid surface, the quick rise in force to a threshold level X will expose a site in the linker protein complex within the leading edge, where SFK is localized and can cause the reinforcement of this site by phosphorylating the linker protein complex. The reinforced site can then stabilize the edge and cause further cell extension. On soft surfaces, however, the force will only rise to X farther away from the edge, and SFKs are not available to phosphorylate the substrate to reinforce the site.

RPTP α -dependent activation of SFK is essential for the rigidity sensing process on FN. In studies of the cellular response to substrate stretch, cytoskeleton stretch activates SFK-dependent phosphorylation of p130Cas (32) and p130Cas^{-/-} cells are defective in matrix rigidity sensing (A. Kostic and M. P. Sheetz, unpublished results). P130Cas is also localized to the leading edge (33). Thus, rigidity sensing appears to involve SFK phosphorylation of substrates at the leading edge.

Models of rigidity sensing

Mechanically, there are two basic mechanisms that could account for the cell's ability to sense matrix rigidity: 1), a time-dependent change in force or 2), a position-dependent change in force. Our data show that rigidity sensing is position dependent and from the characteristics of the laser traps, we can estimate the level of force and the distances involved. With the rapid displacement of the 0.02 pN/nm laser trap by 500 nm, a force of ~ 10 pN is produced that is sufficient to cause increased attachment to the cytoskeleton. Therefore, we suggest that a rapid application of 10 pN of force can elicit the rigidity response. In the case of the rigid trap (0.18 pN/nm), a force of 10–20 pN is reached after 50–100 nm of displacement. Another way of considering the rigidity response is to consider the velocity of actin filament movement in lamellipodia, 60 nm/s (19). At that velocity, the force of 10–20 pN will be reached within 1–2 s. Thus, the rigidity response can be caused either by the cell pulling on a rigid surface or by an active matrix pulling on the cell. In both cases, a rise in force on the cytoskeleton should occur within 50–100 nm of the initial binding site to elicit a rigidity response (Fig. 5).

At a biochemical level, the increase in phosphorylation can be explained by several different mechanisms. One possibility that fits with components known to be involved is a position and force-dependent phosphorylation (Fig. 5). If a phosphorylation site in the FN-cytoskeleton linkage is exposed by force and the kinase that reacts with that site is restricted to the leading edge, then only in the rigid case will

the kinase be close enough to the site to phosphorylate it. Force-dependent activation of p130Cas for SFK phosphorylation has been observed upon cytoskeleton stretch (Y. Sawada, M. Tamada, O. Cherniavskaya, and M. P. Sheetz, unpublished results), and SFKs are able to bind to matrix binding sites. Thus, we believe that FN rigidity sensing occurs at the leading edge of active lamellipodia and produces signals that cause cell migration toward more rigid FN matrices or durotaxis. Further studies are required to define the critical proteins and the actual mechanism of tyrosine phosphorylation in response to substrate rigidity.

We thank G. Giannone, O. Rossier, and H. Xenias for microscope support, and A. Kostic, M. Tamada, A. S. Meshel, H. G. Doebereiner, Y. Sawada, and J. Liu for helpful discussion.

This work was supported by a grant from NIH to M.P.S.

REFERENCES

- Lo, C. M., H. B. Wang, M. Dembo, and Y. L. Wang. 2000. Cell movement is guided by the rigidity of the substrate. *Biophys. J.* 79: 144–152.
- Pelham, R. J. Jr., and Y. Wang. 1997. Cell locomotion and focal adhesions are regulated by substrate flexibility. *Proc. Natl. Acad. Sci. USA.* 94:13661–13665.
- Beningo, K. A., C. M. Lo, and Y. L. Wang. 2002. Flexible polyacrylamide substrata for the analysis of mechanical interactions at cell-substratum adhesions. *Methods Cell Biol.* 69:325–339.
- Cukierman, E., R. Pankov, D. R. Stevens, and K. M. Yamada. 2001. Taking cell-matrix adhesions to the third dimension. *Science.* 294: 1708–1712.
- Georges, P. C., and P. A. Janmey. 2005. Cell type-specific response to growth on soft materials. *J. Appl. Physiol.* 98:1547–1553.
- Yeung, T., P. C. Georges, L. A. Flanagan, B. Marg, M. Ortiz, M. Funaki, N. Zahir, W. Ming, V. Weaver, and P. A. Janmey. 2005. Effects of substrate stiffness on cell morphology, cytoskeletal structure, and adhesion. *Cell Motil. Cytoskeleton.* 60:24–34.
- Sieminski, A. L., R. P. Hebbel, and K. J. Gooch. 2004. The relative magnitudes of endothelial force generation and matrix stiffness modulate capillary morphogenesis in vitro. *Exp. Cell Res.* 297:574–584.
- Engler, A. J., M. A. Griffin, S. Sen, C. G. Bonnemann, H. L. Sweeney, and D. E. Discher. 2004. Myotubes differentiate optimally on substrates with tissue-like stiffness: pathological implications for soft or stiff microenvironments. *J. Cell Biol.* 166:877–887.

9. Semler, E. J., C. S. Ranucci, and P. V. Moghe. 2000. Mechanochemical manipulation of hepatocyte aggregation can selectively induce or repress liver-specific function. *Biotechnol. Bioeng.* 69:359–369.
10. Wang, H. B., M. Dembo, and Y. L. Wang. 2000. Substrate flexibility regulates growth and apoptosis of normal but not transformed cells. *Am. J. Physiol. Cell Physiol.* 279:C1345–C1350.
11. Putman, D. L., D. K. Park, J. S. Rhim, A. F. Steuer, and R. C. Ting. 1977. Correlation of cellular aggregation of transformed cells with their growth in soft agar and tumorigenic potential. *Proc. Soc. Exp. Biol. Med.* 155:487–494.
12. Flanagan, L. A., Y. E. Ju, B. Marg, M. Osterfield, and P. A. Janmey. 2002. Neurite branching on deformable substrates. *Neuroreport.* 13: 2411–2415.
13. Kusaka, K., Y. Harihara, G. Torzilli, K. Kubota, T. Takayama, M. Makuuchi, M. Mori, and S. Omata. 2000. Objective evaluation of liver consistency to estimate hepatic fibrosis and functional reserve for hepatectomy. *J. Am. Coll. Surg.* 191:47–53.
14. Wong, J. Y., A. Velasco, P. Rajagopala, and Q. Pham. 2003. Directed movement of vascular smooth muscle cells on gradient-compliant hydrogels. *Langmuir.* 19:1908–1913.
15. Felsenfeld, D. P., D. Choquet, and M. P. Sheetz. 1996. Ligand binding regulates the directed movement of beta1 integrins on fibroblasts. *Nature.* 383:438–440.
16. Schwarz, U. S., N. Q. Balaban, D. Riveline, A. Bershadsky, B. Geiger, and S. A. Safran. 2002. Calculation of forces at focal adhesions from elastic substrate data: the effect of localized force and the need for regularization. *Biophys. J.* 83:1380–1394.
17. Mandeville, J. T., and F. R. Maxfield. 1997. Effects of buffering intracellular free calcium on neutrophil migration through three-dimensional matrices. *J. Cell. Physiol.* 171:168–178.
18. Giannone, G., B. J. Dubin-Thaler, H. G. Dobreiner, N. Kieffer, A. R. Bresnick, and M. P. Sheetz. 2004. Periodic lamellipodial contractions correlate with rearward actin waves. *Cell.* 116:431–443.
19. Aznavoorian, S., M. L. Stracke, J. Parsons, J. McClanahan, and L. A. Liotta. 1996. Integrin alphavbeta3 mediates chemotactic and haptotactic motility in human melanoma cells through different signaling pathways. *J. Biol. Chem.* 271:3247–3254.
20. Li, X., B. Chen, S. D. Blystone, K. P. McHugh, F. P. Ross, and D. M. Ramos. 1998. Differential expression of alphav integrins in K1735 melanoma cells. *Invasion Metastasis.* 18:1–14.
21. Kiosses, W. B., S. J. Shattil, N. Pampori, and M. A. Schwartz. 2001. Rac recruits high-affinity integrin alphavbeta3 to lamellipodia in endothelial cell migration. *Nat. Cell Biol.* 3:316–320.
22. von Wichert, G., G. Jiang, A. Kostic, K. De Vos, J. Sap, and M. P. Sheetz. 2003. RPTP-alpha acts as a transducer of mechanical force on alphav/beta3-integrin-cytoskeleton linkages. *J. Cell Biol.* 161: 143–153.
23. Ponniah, S., D. Z. Wang, K. L. Lim, and C. J. Pallen. 1999. Targeted disruption of the tyrosine phosphatase PTPalpha leads to constitutive downregulation of the kinases Src and Fyn. *Curr. Biol.* 9:535–538.
24. Su, J., M. Muranjan, and J. Sap. 1999. Receptor protein tyrosine phosphatase alpha activates Src-family kinases and controls integrin-mediated responses in fibroblasts. *Curr. Biol.* 9:505–511.
25. Petrone, A., F. Battaglia, C. Wang, A. Dusa, J. Su, D. Zagzag, R. Bianchi, P. Casaccia-Bonnel, O. Arancio, and J. Sap. 2003. Receptor protein tyrosine phosphatase alpha is essential for hippocampal neuronal migration and long-term potentiation. *EMBO J.* 22: 4121–4131.
26. Jiang, G., G. Giannone, D. R. Critchley, E. Fukumoto, and M. P. Sheetz. 2003. Two-piconewton slip bond between fibronectin and the cytoskeleton depends on talin. *Nature.* 424:334–337.
27. Chen, B. H., J. T. Tzen, A. R. Bresnick, and H. C. Chen. 2002. Roles of Rho-associated kinase and myosin light chain kinase in morphological and migratory defects of focal adhesion kinase-null cells. *J. Biol. Chem.* 277:33857–33863.
28. Avizienyte, E., A. W. Wyke, R. J. Jones, G. W. McLean, M. A. Westhoff, V. G. Brunton, and M. C. Frame. 2002. Src-induced deregulation of E-cadherin in colon cancer cells requires integrin signalling. *Nat. Cell Biol.* 4:632–638.
29. Levinson, H., J. E. Hopper, and H. P. Ehrlich. 2002. Overexpression of integrin alphav promotes human osteosarcoma cell populated collagen lattice contraction and cell migration. *J. Cell. Physiol.* 193: 219–224.
30. Kaido, T., M. Yebra, V. Cirulli, and A. M. Montgomery. 2004. Regulation of human beta-cell adhesion, motility, and insulin secretion by collagen IV and its receptor alpha1beta1. *J. Biol. Chem.* 279:53762–53769.
31. Giannone, G., G. Jiang, D. H. Sutton, D. R. Critchley, and M. P. Sheetz. 2003. Talin1 is critical for force-dependent reinforcement of initial integrin-cytoskeleton bonds but not tyrosine kinase activation. *J. Cell Biol.* 163:409–419.
32. Tamada, M., M. P. Sheetz, and Y. Sawada. 2004. Activation of a signaling cascade by cytoskeleton stretch. *Dev. Cell.* 7:709–718.
33. Abassi, Y. A., M. Rehn, N. Ekman, K. Alitalo, and K. Vuori. 2003. p130Cas couples the tyrosine kinase Bmx/Etk with regulation of the actin cytoskeleton and cell migration. *J. Biol. Chem.* 278:35636–35643.